



**HAL**  
open science

## Wood excavation, construction, and architecture in two Reticulitermes subterranean termites

L. Berville, E. Darrouzet

► **To cite this version:**

L. Berville, E. Darrouzet. Wood excavation, construction, and architecture in two Reticulitermes subterranean termites. *Insectes Sociaux*, 2019, 66 (3), pp.403 - 411. 10.1007/s00040-019-00696-x . hal-02794520

**HAL Id: hal-02794520**

**<https://univ-tours.hal.science/hal-02794520>**

Submitted on 19 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Click here to view linked References](#)Wood excavation, construction, and architecture in two *Reticulitermes* subterranean termites

Laurence Berville, Eric Darrouzet\*

IRBI, UMR CNRS 7261, University of Tours, Parc de Grandmont, 37200 Tours, France

\*Corresponding author: eric.darrouzet@univ-tours.fr

**Abstract:**

Collective constructions are marvels of complexity, composed of networks of tunnels and chambers. However, it is difficult to study subterranean nests without using invasive techniques because the nests are built within pieces of wood and/or in the soil. Using computerized tomography scans and medical imaging software (OsiriX), we were able to observe nest creation, constructions, and architecture of two subterranean termite species. We monitor the nests' growth in three dimensions built by two *Reticulitermes* species: *R. grassei*, a species native to Europe, and *R. flavipes*, an invasive species introduced from North America, over a several-month period. Doing so, we wanted to know whether the construction of the nest could participate to the invasive success of *R. flavipes*. Although the two species displayed some similarities (*i.e.*, nest creation, chamber size, and levels of wood consumption), only *R. flavipes* built interior structures. Some of these structures changed over time and thus might play a role in the trade-off between wood consumption, colony protection, and environmental homeostasis.

**Significance Statement**

Nest architectures and their management is one of the triumphs of social animals such as social insects. The functional analysis of those structures composed of networks of tunnels and chambers could help to understand the biology and ecology of their builders. Excavation, consumption and nest design are important parameters in explaining a number of interesting biological traits such as the establishment of invasive species. Some recent studies have introduced X-ray tomography as a tool to analyze insect architectures. We employed this non-invasive technique to observe nest creations, construction and evolution in time, and resolve three-dimensional nest architecture. We analyzed a native and an invasive subterranean termite species. Interestingly, only the invasive species built specific interior structures which change over time. Our study leads to a hypothesis of optimal chamber number, which balances colony wood consumption against nest functionality. The constructions could play a role for environmental homeostasis that seems not necessary for the native species.

**Keywords:** animal architecture, tomography, *Reticulitermes flavipes*, *R. grassei*, self-organized structures, social insects

## 36 Introduction

37 Animals can create controlled microhabitats using self-made structures. The most impressive are the nests of  
38 social insects (i.e., ants, bees, wasps, and termites). Some termite mounds rise higher than 10 meters, which is  
39 more than 1,000 times the height of a worker. Even more intriguing is that such nests are produced without  
40 individual workers using any sort of collective “blueprint” (Buhl et al. 2005; Deneubourg and Franks, 1995).  
41 Social insects have achieved remarkable ecological success and dominance (Hölldobler and Wilson, 1990),  
42 perhaps due in part to their nests, which provide benefits in terms of defense, shelter (Traniello and Leuthold,  
43 2000), controlled microclimatic conditions (e.g., air composition [ants: Bollazzi et al. 2012], temperature and  
44 humidity [termites: Ruelle 1964]), structure, and, in some cases, food. Indeed, certain subterranean termite  
45 species also consume the wood in which they live (Grassé 1984). These species serve as excellent models for  
46 studying nest development as they must trade-off between food consumption (wood) and environmental  
47 homeostasis. Part of this process is controlling the number and size of both tunnels and chambers. Thus, nest  
48 architecture may play an important role in regulating colony activity.

49 However, subterranean lifestyle makes it difficult to characterize nest structure. Previous studies, **which**  
50 **analyzed nest architectures in several insect species**, have largely employed destructive methods such as  
51 fragmentation (Bouillon 1964), step-by-step excavation (King and Spink 1969; Rabeling et al. 2007; Tschinkel  
52 1999), the tracing of radioisotope-labeled foragers (Spragg and Paton 1980), or casts (i.e., latex [Garner 1953],  
53 cement [Bollazzi et al. 2012; Ruelle 1964], orthodontic plaster, wax, aluminum, zinc, or lead [Tschinkel 2010,  
54 2011]). However, these approaches can only yield a temporal and/or spatial snapshot of the nest. They cannot  
55 provide information about structural dynamics, such as the growth of tunnel networks, nor can they be used to  
56 follow nest creation, wood consumption, and/or building activity over time. More recent studies, though, have  
57 begun utilizing computerized tomography (CT) scans to non-invasively visualize and characterize the  
58 architecture of social insect nests (ants [Halley et al. 2005; Khuong et al. 2016; Minter et al. 2012; **Monaenkova**  
59 **et al. 2015**]; bees [Greco et al. 2006]; termites [Darrouzet 2013; Eom et al. 2015; Fuchs et al. 2004; Himmi et al.  
60 2014, 2016; Perna et al. 2008a,b]; wasps [Darrouzet 2013]; and hornets [Darrouzet 2013; Rome et al. 2015]).  
61 Using CT scans, it is possible to reconstruct a 3D structure via a series of 2D images; the structure’s inhabitants  
62 are unharmed, even after several scans. Thanks to the images’ high-contrast resolution, structural differences in  
63 physical density can be distinguished (see [Sup. file Video 1](#)). The ability of structures to block the X-ray beam  
64 can also be characterized. Nevertheless, despite their potential, longitudinal studies using CT scans have only  
65 been carried in ants (*Linepithema humile*: 7 or 8 scans over 6 months [Halley et al. 2005] and *Lasius flavus*: 8  
66 scans over 48 h [Minter et al. 2012]). Their use may be tempered by the tool’s main limitation: scanner size. It is  
67 impossible to analyze large structures.

68 **An important requirement for the management of invasive species is to identify biological and**  
69 **ecological factors that influence their ability to establish and spread within a new environment, and how they**  
70 **present a significant advantage against native species. For example, *Reticulitermes flavipes* and *R. grassei* are**  
71 **subterranean termite species which live in sympatry along the French Atlantic coast, where their colonies exhibit**  
72 **important differences in their social organization, behavior, and breeding system (Perdereau et al. 2010, 2011,**  
73 **2015). *R. grassei* is native to southwestern Europe (France and Spain). *R. flavipes*, in contrast, is an invasive**  
74 **species that is native to the eastern United States but that has spread to South America (Aber and Fontes 1993;**  
75 **Su et al. 2006) and Europe (Dronnet et al. 2006; see Evans 2013 for a review). Both species belong to the family**  
76 **Rhinotermitidae and, as such, share certain traits. Some studies demonstrated that *R. flavipes* has some**  
77 **advantages against *R. grassei*. For example, interspecific competition showed that *R. flavipes* was dominant over**  
78 ***R. grassei*, and this invasive species foraged over a greater distance than *R. grassei* colonies (Perdereau et al,**

79 2011). Consequently, this competitive asymmetry may enable *R. flavipes* to become dominant in the  
80 environment. Interestingly, the main difference between the two species concerns their capacity to produce  
81 secondary neotenic reproductives. Genetic studies revealed that all French *R. flavipes* colonies produce more  
82 neotenic than *R. grassei* colonies (Leniaud et al, 2011). In such conditions, its interspecific superiority, its lack  
83 of intraspecific aggression, the production of numerous neotenic and large extensive colonies, seem to be some  
84 of the reasons for *R. flavipes* invasive success. Among these factors, nest-building dynamics, nest shapes and  
85 sizes, wood consumption dynamics are largely unknown for both species. However, information on nest  
86 dynamics is essential to understand the wood consumption dynamic and to assess the efficacy of the strategies  
87 used to control invasive termite infestations (Forschler 1994). For example, they both eat and live in non-  
88 decomposed wood. They are initially single-piece nesters, but they eventually move to a new piece of wood  
89 when their original nests have been completely consumed (Evans 2013). Both produce large subterranean  
90 colonies and are therefore ideal species for studying wood consumption.

91 In this study, we sought to discover if both *Reticulitermes* species build interior structures and thereby  
92 to discover the relationship between these constructions and the volume and dynamic of wood consumed. Doing  
93 so, we wanted to know whether the construction of the nest could participate to the invasive success of *R.*  
94 *flavipes*. As such species seems to produce larger colonies and numerous neotenic, we supposed that (1) wood  
95 consumption would be greater in the invasive species (to feed numerous individuals and more particularly  
96 neotenic), (2) *R. flavipes* workers would have greater capacities to build the nest, and consequently (3) nest  
97 structures from both species could present some differences in size and shape. Moreover, our study leads to a  
98 hypothesis of optimal chamber number, which balances colony wood consumption against nest functionality.  
99 We hypothesis that a cumulative Weibull distribution or a sigmoid function will appropriately models the  
100 biological process at work.

101 To test these hypotheses, we used CT scans to characterize nest creation, construction, and architecture  
102 as well as wood consumption over a ten-month period in experimental fragment colonies of both species in the  
103 same experimental conditions. More specifically, we exploited CT scans to investigate (1) nest construction  
104 dynamics, (2) wood consumption, and (3) intercolonial variation in both.

105

## 106 **Materials and Methods**

### 107 *Colony rearing conditions*

108 *Reticulitermes* colonies were collected from wood fragments and pine tree stumps in the St. Trojan  
109 forest on Oléron Island (Charente-Maritime, France). They were identified to species using cuticular  
110 hydrocarbon profiles (Bagnères et al. 1990, 1991). For the duration of the experiment, colonies were kept at  
111 26 °C in full darkness. We then filled plastic boxes (18 x 12 x 7 cm) with moist Fontainebleau sand (very fine,  
112 homogeneous granularity); sand depth was 2 mm. We added a block of untreated pine (12 x 4 x 4 cm) as a food  
113 source. The sand was moistened every day. Four fragment colonies of *R. grassei* and six fragment colonies of *R.*  
114 *flavipes* were introduced into separate boxes; each contained 900 workers and was taken from a different field-  
115 collected colony.

116

### 117 *Computerized X-ray tomography scan analysis*

118 To monitor wood consumption, each plastic box was subject to CT scans (120 keV, 180 mAs) using a  
119 Phillips Brilliance CT 40 medical scanner, made available to us by the Neuroradiology Department at the Tours  
120 Hospital (Tours, France). We obtained a series of virtual vertical “slices” through the box (thickness of 0.67 mm,  
121 increments of 0.33 mm) (Fig. 1a and [Sup file 1](#)). The boxes were scanned before the termites were introduced

122 (T0) and then at regular intervals post introduction (Sup file 1): at 13 days (T1), 20 days (T2), 51 days (T3), 65  
123 days (T4), 100 days (T5), 202 days (T6, Sup file video 1), and 258 days (T7).

124 Then, we analyzed each series of CT scans (DICOM standard; Fig. 1a, Sup file video 1) using OsiriX v.  
125 4.19 image processing software (Darrouzet, 2013; Rosset et al. 2004), which is designed to allow 3D volumes to  
126 be reconstructed from large sets of multidimensional and multimodality images (Fig. 1b,c and Sup file video 2).  
127 Areas of wood consumption, which we called “regions of interest” (ROIs), were identified manually on each  
128 image and measured (Fig. 1a). We thus obtained precise information about ROI size, shape, and boundaries. We  
129 were also able to observe nest architecture over time, namely changes in tunnels (interior and exterior), chamber  
130 number, wood consumption volume, and interior structures. One of the *R. grassei* colonies died between T5 and  
131 T6, but we used the data available from T0 to T5.

132

### 133 *Statistics*

134 Our statistical analyses were performed using SigmaStat (v. 10.0.054). In all cases, data normality was  
135 first assessed using Kolmogorov-Smirnov tests. In the results, means are presented with standard errors. To  
136 compare the locations where the two species initially attacked the wood pieces, t-tests (normally distributed data)  
137 and Mann-Whitney rank sum tests (non-normally distributed data) were used. To assess differences in chamber  
138 volume and number across time for the two species, two-way repeated measures ANOVAs was used; *a*  
139 *posteriori* pairwise comparisons were performed using Holm-Sidak tests.

140 To test if the data is suitable for a regression analysis we used a Kolmogorov-Smirnov test for a  
141 normally distributed population and a constant variance test by computing the Spearman rank correlation  
142 between the absolute values of the residuals and the observed value of the dependent variable. To gauge the  
143 ability of the regression equation to predict the dependent variable, an ANOVA (F test) was performed. To  
144 provide a measure of how well the number of chamber is dependent of the volume consumed, the coefficient of  
145 determination ( $r^2$ ) was used.

146

## 147 **Results**

### 148 *Nest creation*

149 During the nest-creation period (T1 and T2), there was no clear pattern in the way the termites attacked  
150 the wood (Mann-Whitney U Statistic,  $P=0.33$ ): they targeted both the bottom face of the block (*R. grassei*,  $n=50$ ;  
151 *R. flavipes*,  $n=34$ ) and the exposed faces of the block (*R. grassei*,  $n=50$ ; *R. flavipes*,  $n=14$ ). The first exterior  
152 tunnels, made of sand, wood, and fecal pellets, were observed after 51 days; these afforded protection to the  
153 termites. From that point on, it was rare to observe an attack that was not associated with a sand tunnel (*R.*  
154 *grassei*,  $n=9$ ; *R. flavipes*,  $n=6$ ), as most of the new chambers were initiated from inside the tunnels (*R. grassei*,  
155  $n=215$ ; *R. flavipes*,  $n=684$ ), or via the extension of other chambers (*R. grassei*,  $n=188$ ; *R. flavipes*,  $n=253$ ). The  
156 two species did not differ in their overall approach (T3 and T7;  $t= -0.994$ ;  $P=0.359$ ) (Fig. 2).

157

### 158 *Wood consumption*

159 As they ate through the wood, the termites generated living space for themselves. We identified 1,829  
160 ROIs across the 10 study colonies; their mean size was  $0.43 \text{ cm}^3$  (range:  $0.0004 - 10.25 \text{ cm}^3$ ). Most ROIs were  
161 small ( $50\% < 0.111 \text{ cm}^3$  and  $90\% < 1.196 \text{ cm}^3$ ) (Fig. 3).

162 At each time point, the two termite species had equivalent chamber numbers (two-way repeated  
163 measures ANOVA:  $F=1.287$ ,  $df=1$ ,  $P=0.289$ ) after accounting for differences in the factor times ( $F=0.648$ ,  $df=7$ ,  
164  $P=0.714$ ). Both species appeared to go through three phases of wood consumption (Fig. 4;  $F=37,945$ ,  $df=7$ ,

165 **P<0.001**): there were no differences in chamber number at T0, T1, and T2 (phase one), at T3 and T4 (phase two),  
 166 and at T5, T6, and T7 (phase three). During the first phase, a small number of new chambers were made. During  
 167 the second phase, chamber creation accelerated. During the third phase, chamber creation declined (Fig. 4). By  
 168 the end of the experiment, both species had generated almost the same number of chambers. *R. flavipes* nests  
 169 contained between 33 and 64 chambers, while *R. grassei* nests contained between 24 and 53 chambers (Fig. 4).

170 There also did not appear to be interspecific differences in wood consumption over time (T0 to T7; two-  
 171 way repeated measures ANOVA: **F= 0.774**, P=0.404). At the end of the experiment (*i.e.*, after 258 days), *R.*  
 172 *flavipes* colonies had eaten 15.10 – 43.66 cm<sup>3</sup> of wood, corresponding to a mean 15.51% ± 2.068 of **total wood**  
 173 **block volume** (Fig. 4). *R. grassei* had eaten 21.19 – 32.26 cm<sup>3</sup> of wood, corresponding to a mean 14.2% ± 1.83  
 174 of total wood block volume (Fig. 4). While the volume of wood consumed was equivalent between species at  
 175 each time point, the mean consumption across time for species was different (two-way repeated measures  
 176 ANOVA: **F=73,347**, P<0.001). Within species, all pairwise comparisons between time points were statistically  
 177 significant, except during the first stage (T0 vs. T1, T0 vs. T2, and T1 vs. T2) and for T3 versus T4 (Fig. 4).  
 178 Chamber number was related to the percentage of wood volume consumed in both species (*R. flavipes*: r<sup>2</sup>=0.979,  
 179 F=65.07, P=0.0008; *R. grassei*: r<sup>2</sup>=0.984, F=84.27, P=0.0005; Fig. 5). The best fit for the data was a four-  
 180 parameter non-linear regression model with f(x) = y<sub>0</sub>+a/(1+exp<sup>(-(x-x<sub>0</sub>)/b)</sup>). The equations were as follows:

$$R. grassei \quad y = -2.2 + \frac{39.80}{\left(1 + e^{\left(\frac{-(x-3.85)}{1.24}\right)}\right)}$$

$$R. flavipes \quad y = -0.22 + \frac{47.98}{\left(1 - e^{\left(\frac{-(x-3.85)}{1.72}\right)}\right)}$$

181

182

183

184 with x = percentage of wood volume consumed, y= chamber number, a = the minimum value that can be  
 185 obtained (*i.e.* when x=0) and b = Hill's slope of the curve. An Anova was used accordingly, demonstrating that  
 186 the volumes of wood consumed by both species contribute to the prediction of the number of chamber observed  
 187 in the nests.

188

189

#### 190 *Interior structures*

191 Only *R. flavipes* seemed to have the ability to generate and modify interior architectural structures such  
 192 as pillars, ceilings/floors, tunnels, and inner/outer walls (Fig. 2). The timing was colony specific: the first  
 193 example of such structures occurred after 20 days in one colony (Fig. 8), but after 258 days in another colony.  
 194 By the end of the experiment, a mean of 14.5 ± 7 interior structures were present in *R. flavipes* nests. Although  
 195 there was marked intercolonial variability, there was a strong correlation between the number of days that had  
 196 passed and the number of such structures (r<sup>2</sup>=0.992, Anova F=747.39, P=0.0001; Fig. 8). There was a significant  
 197 relationship between the number of interior structures and the number of chambers (r<sup>2</sup>=0.974; F=50.45;  
 198 P=0.0012; Fig. 9). The best fit for the data was a four-parameter equation (Weibull distribution), where x=the  
 199 number of constructions:

$$y = 47.27 \left(1 - e^{\left(\frac{-(x + 70.4)}{22.29}\right)^{10.03}}\right)$$

200

201

202 **Discussion**

203 This study used CT scans (X-ray tomography) to provide the first detailed quantitative descriptions of  
204 *Reticulitermes* nests **dynamics**. We obtained new information about nest creation, the dynamics of wood  
205 consumption, the creation of chambers and tunnel networks, and shifts in interior architecture. Because initial  
206 conditions were standardized (i.e., wood block size and colony size), we were able to characterize and compare  
207 these features over time.

208 More specifically, we collected data on two sympatric *Reticulitermes* species, one native and one  
209 invasive, which has helped enhance our understanding of their natural histories. First, we saw no interspecific  
210 differences in the volume of wood consumed or in the number of chambers created. Second, we observed three  
211 phases of wood consumption. Over the first 25 days of the experiment (Phase 1), wood consumption was low.  
212 Then, the first exterior tunnels appeared, wood consumption accelerated, and a large number of small chambers  
213 were created (Phase 2). Finally, 200 days into the experiment, the speed of wood consumption remained  
214 unchanged, but a slightly smaller number of chambers were generated. However, these chambers were either  
215 larger or the product of fusion.

216 By eating wood and engaging in building activity over a several-month period, termites can create  
217 networks of interconnected galleries and chambers in which they live and through which they can travel (Grassé  
218 1984, also see [Sup file video 1 and 2](#)). Within the nest, termites are also protected from predators and climatic  
219 extremes. Nonetheless, a major challenge remains: sufficient ventilation (Ohashi et al. 2012). Air quality within  
220 the nest can be compromised by colony respiration, namely the production and release of CO<sub>2</sub> (Risch et al.  
221 2012). In general, ventilation systems are passive, driven by temperature or velocity gradients in air currents  
222 (Wenzel 1990). Indeed, termites are well known for their ability to regulate nest conditions, including  
223 temperature and humidity (Lüscher 1955; Korb and Linsenmair 2000; Korb 2003; see also Turner 2001 for a  
224 review). The construction of different interior structures could help enhance air flow and thus improve air  
225 quality. To this end, nest configurations might need to remain dynamic. The small size of most of the chambers  
226 observed in this study could be the product of a trade-off between wood consumption, colony defense, and  
227 environmental homeostasis. As colonies grow, old chambers can be rearranged and new chambers can be  
228 created: here, we observed that some chambers increased in size or fused.

229 One of our notable results is that, although both species displayed similar levels of wood consumption  
230 and created exterior tunnels, only *R. flavipes* built interior structures. These included pillars, ceilings/floors, and  
231 walls (Fig. 1b). Like *Cubitermes* species (Perna et al. 2008a), *R. flavipes* could adjust interior nest architecture  
232 by adding and modifying the number of interior structures, depending on chamber number or wood  
233 consumption. Ceilings and floors appeared in the largest chambers, and some old entrances were sealed off.  
234 Since colonies were given just one piece of wood, termites were forced to eat where they lived. Consequently,  
235 for *R. flavipes*, such structures may be part of the trade-off between wood consumption, colony defense, and  
236 environmental homeostasis. In our experiment, however, exterior temperature and humidity were kept constant,  
237 and the termites were enclosed in boxes, preventing any air movement. Although both species were collected  
238 from the same area and thus experienced the same environmental conditions, it could be that *R. grassei*, the  
239 native species, did not build interior structures because laboratory conditions were acceptable. Perhaps this was  
240 not the case for *R. flavipes*, the invasive species, which may have needed to manage nest ventilation. Perna et al.  
241 (2008a) showed that, in *Cubitermes* species, final nest topology reflects a compromise between efficiency and  
242 defense. The same could be true in *Reticulitermes* species. Furthermore, it could be that *R. flavipes* utilizes  
243 interior structures to help defend against predators in its native range within North America and that this

244 behavior was conserved within its introduced range in France. However, *R. flavipes*' potential natural predators  
245 remain unidentified.

246 Another hypothesis for the interspecific difference in the use of interior structures could be linked to the  
247 production of neotenics. More specifically, *R. flavipes* may produce more neotenics than *R. grassei*: Leniaud et  
248 al. (2011) observed that, after one year, 100% of *R. flavipes* groups (composed of workers) had produced  
249 neotenics, while just 63% of *R. grassei* groups had done the same. Compared to *R. grassei*, *R. flavipes* produced  
250 significantly more offspring because female numbers and productivity were both higher. Here, consequently, our  
251 *R. flavipes* colonies may have been more densely populated than our *R. grassei* colonies after just a few months.  
252 Higher densities could have led to higher CO<sub>2</sub> concentrations and thus triggered the implementation of a  
253 ventilation system mediated via interior structures. It is also conceivable that *R. flavipes* neotenics are more  
254 sensitive to certain environmental conditions, requiring more intensive nest management. Unfortunately, because  
255 we did not assess the characteristics of our colonies at the end of the study, additional research is needed to test  
256 these hypotheses.

257 In our study, there was noticeable intercolonial variation in nest configuration (Sup File 1), even though  
258 nests presumably share the same functional purpose and both colony numbers and wood consumption levels  
259 were similar. There were no interspecific differences in chamber organization. Nest initiation was also similar.  
260 Although the colonies attacked the wood block at different locations, all their entrance cavities were excavated in  
261 the springwood and the network expanded along the wood's microstructure. This pattern is evident on CT scans  
262 (Fig 1c) because the degree of X-ray absorption is correlated with wood density, which is greatest for the annual  
263 growth rings (Himmi et al. 2016). We found that both termite species generally excavated tunnels and chambers  
264 in softer wood (Sup file 2). This particular behavior is known to be affected by wood chemistry (Abe and  
265 Higashi 1991) and nutritional value (Shellman-Reeve 1994). Only a few tunnels crossed harder wood to reach  
266 softer areas, which allowed nest development.

267 Taken together, our results suggest that *R. grassei* and *R. flavipes* differ in their nest-building design.  
268 This study confirms that both the environment and the termites themselves can influence the expression of  
269 extended phenotypes, such as nest configuration. Although substrate quality (e.g., winter/summer wood) can  
270 influence nest architecture making them nest specific, while, building behavior is species specific. This study  
271 also underscores the utility of CT scans as a non-invasive technique for following wood consumption and nest  
272 architecture in termites over time and the expression of extended phenotypes. Indeed, the types of data it yields  
273 could be used in future research efforts to model nest construction dynamics.

274

## 275 **Acknowledgments**

276 We gratefully acknowledge J. Pearce for her English revisions. We would like to thank S. Dupont for  
277 rearing the termite colonies in the laboratory and D. Herbreteau for letting us use the CT scanner at Tours  
278 Hospital.

279

## 280 **References**

- 281 Abe T, Higashi M (1991) Cellulose centred perspective on terrestrial community structure. *Oikos* 60:127–133  
282 Aber A, Fontes LR (1993). *Reticulitermes lucifugus* (Isoptera, Rhinotermitidae), a pest of wooden structures, is  
283 introduced into the South American Continent. *Sociobiology* 21:335-339.  
284 Bagnères A-G, Clément J-L, Blum M, Severson R, Joulie C, Lange C (1990). Cuticular hydrocarbons and  
285 defensive compounds of *Reticulitermes flavipes* (Kollar) and *R. santonensis* (Feytaud): polymorphism  
286 and chemotaxonomy. *J Chem Ecol* 16:3213–3244



287 Bagnères A-G, Killian A, Clément J-L, Lange C (1991). Interspecific recognition among Termites of the genus  
288 *Reticulitermes*. Evidence for a role for the cuticular hydrocarbons. *J Chem Ecol* 17:2397-2420

289 Bollazzi M, Forti LC, Roces F (2012). Ventilation of the giant nests of *Atta* leaf-cutting ants: Does underground  
290 circulating air enter the fungus chambers? *Insectes Soc.* 59(4):487–498

291 Bouillon A (1964). Structure et accroissement des nids d'*Apicotermes Holmgren* (Isoptere, Termitinae). 295-  
292 326pp. In: Etudes sur les termites Africains. A. Bouillon (Eds), Université de Léopoldville.

293 Buhl J, Deneubourg JL, Grimal A, Theraulaz G (2005). Self-organized digging activity in ant colonies. *Behav*  
294 *Ecol and Socio* 58(1):9–17

295 Darrouzet E (2013). Les insectes bâtisseurs : nids de termites, de guêpes et de frelons. *Connaissances & Savoirs*  
296 (Eds).

297 Deneubourg JL, Franks NR (1995). Collective control without explicit coding: The case of communal nest  
298 excavation. *J Insect Behav* 8(4):417–432

299 Dronnet S, Lohou C, Christidès J-P, Bagnères A-G (2006). Cuticular hydrocarbon composition reflects genetic  
300 relationship among colonies of the introduced termite *Reticulitermes santonensis* Feytaud. *J Chem Ecol*  
301 30:1027–1042

302 Eom Y-H, Perna A, Fortunato S, Darrouzet E, Theraulaz G, Jost C (2015). Network based model of the growth  
303 of termite nests. *Phy Res E* 92(6): 062810

304 Evans TA, Forschler BT, Kenneth G (2013). Biology of invasive Termites: A worldwide review. *Annu Rev*  
305 *Entomol* 58:455-74

306 Forschler BT (1994), Fluorescent spray paint as a topical marker on subterranean termites (Isoptera:  
307 *Rhinotermitidae*). *Sociobiology*, 24, pp. 27–38

308 Fuchs A, Schreyer A, Feuerbach S, Korb J (2004). A new technique for termite monitoring using computer  
309 tomography and endoscopy. *Inter J Pest Manag* 50(1):63–66

310 Garner M R (1953). The Preparation of Latex Casts of Soil Cavities for the Study of Tunneling Activities of  
311 Animals. *Science*, 118(3066):380–381

312 Grassé PP (1984). *Termitologia*. Fondation des sociétés - Construction. Tome2. Paris: Masson.

313 Greco M, Bell M, Spooner-Hart R, Holford P (2006). X-ray computerized tomography as a new method for  
314 monitoring *Amegilla holmesi* nest structures, nesting behaviour, and adult female activity. *Entomol Exp*  
315 *et App* 120(1):71–76

316 Halley JD, Burd M, Wells P (2005). Excavation and architecture of Argentine ant nests. *Insectes Soc.* 52(4):350–  
317 356

318 Himmi SK, Yoshimura T, Yanase Y, Oya M, Torigoe T, Imazu S (2014). X-ray tomographic analysis of the  
319 initial structure of the royal chamber and the nest-founding behavior of the drywood termite *Incisitermes*  
320 *minor*. *J Wood Science* 60(6):453–460

321 Himmi SK, Yoshimura T, Yanase Y, Mori T, Torigoe T, Imazu S (2016). Wood anatomical selectivity of  
322 drywood termite in the nest-gallery establishment revealed by X-ray tomography. *Wood Science Tech*  
323 50(3):631–643

324 Hölldobler B, Wilson E (1990). Colony Odor and Kin Recognition. In *The Ants* (pp. 197–199).

325 Khuong A, Gautrais J, Perna A, Sbaï C, Combe M, Theraulaz G (2016). Stigmergic construction and  
326 topochemical information shape ant nest architecture. *PNAS* 113(5):201509829

327 King EG, Spink WT (1969). Foraging Galleries of the Formosan Subterranean Termite, *Coptotermes*  
328 *formosanus*, in Louisiana. *An Entomol Soc Am* 62(3):536–542

329 Korb J, Linsenmair KE (2000) Thermoregulation of termite mounds: what role does ambient temperature and  
330 metabolism of the colony play? *Insectes Soc.* 47:357-363

331 Korb J (2003). Thermoregulation and ventilation of termite mounds. *Naturwissenschaften* 90:212-219

332 Leniaud L, Darrouzet E, Dedeine F, Ahn K, Huang Z, Bagnères A-G (2011). Ontogenic potentialities of the  
333 worker caste in two sympatric subterranean termites in France. *Evol Dev* 13:138–148

334 Lüscher M, (1955). Der Sauerstoffverbrauch bei Termiten und die Ventilation des Nestes bei *Macrotermes*  
335 *natalensis* (Haviland). *Acta Trop* 12:289-307

336 Minter NJ, Franks NR, Robson Brown KA (2012). Morphogenesis of an extended phenotype: four-dimensional  
337 ant nest architecture. *J Royal Soc Interface* 9(68):586–595

338 **Monaenkova D, Gravish N, Rodriguez G, Kutner R, Goodisman MD, Goldman, DI. (2015). Behavioral and**  
339 **mechanical determinants of collective subsurface nest excavation. *J. Exp. Biol.* 218, 1295-1305.**  
340 **doi:10.1242/jeb.113795.**

341 Ohashi M, Domisch T, Finér L, Jurgensen MF, Sundström L, Kilpeläinen J, Risch AC, Niemelä P (2012). The  
342 effect of stand age on CO<sub>2</sub> efflux from wood ant (*Formica rufa* group) mounds in boreal forests. *Soil*  
343 *Biol. Biochem* 52:21–28

344 Perna A, Jost C, Couturier E, Valverde S, Douady S, Theraulaz G (2008a). The structure of gallery networks in  
345 the nests of termite *Cubitermes* spp. revealed by X-ray tomography. *Naturwissenschaften* 95(9) :877–884

346 Perna A, Valverde S, Gautrais J, Jost C, Solé R, Kuntz P, Theraulaz G (2008b). Topological efficiency in three-  
347 dimensional gallery networks of termite nests. *Physica A: Stat Mech App* 387(24):6235–6244

348 **Perdereau E, Bagnères AG, Dupont S, Dedeine F. (2010). High occurrence of colony fusion in a European**  
349 **population of the American termite *Reticulitermes flavipes*. *Insectes sociaux*, 57(4), 393-402.**

350 **Perdereau E, Dedeine F, Christidès JP, Dupont S, Bagnères AG. (2011). Competition between invasive and**  
351 **indigenous species: an insular case study of subterranean termites. *Biological Invasions*, 13(6), 1457-**  
352 **1470.**

353 **Perdereau E, Bagnères AG, Vargo EL, Baudouin G, Xu Y, Labadie P, Dupont S, Dedeine F. (2015).**  
354 **Relationship between invasion success and colony breeding structure in a subterranean termite. *Molecular***  
355 ***ecology*, 24(9), 2125-2142.**

356 Rabeling C, Verhaagh M, Engels W (2007). Comparative study of nest architecture and colony structure of the  
357 fungus-growing ants, *Mycocepurus goeldii* and *M. smithii*. *J Insect Sci* 7(40):1–13

358 Risch AC, Anderson TM, Schütz M (2012). Soil CO<sub>2</sub> emissions associated with termitaria in Tropical savanna:  
359 evidence of hot-spot compensation. *Ecosystems* 15:1147–1157

360 Rome Q, Muller FJ, Touret-Alby A, Darrouzet E, Perrard A, Villemant C (2015). Caste differentiation and  
361 seasonal changes in *Vespa velutina* (Hym.: Vespidae) colonies in its introduced range. *J App Entomol*  
362 139(10):771–782

363 Rosset A, Spadola L, Ratib OJ (2004). OsiriX: An Open-Source Software for Navigating in Multidimensional  
364 DICOM Images. *Digit Imaging* 17:205

365 Ruelle J (1964). L'Architecture du nid de *Macrotermes natalensis* et son sens fonctionnel. In A. Bouillon (Ed.),  
366 Etudes sur les termites Africains. Université de Léopoldville.

367 Shellman-Reeve JS (1994) Limited nutrients in a dampwood termite: nest preference, competition and  
368 cooperative nest defence. *J Anim Ecol* 63:921–932

369 Spragg WT, Paton R (1980). Tracing, Trophallaxis and Population Measurement of Colonies of Subterranean  
370 Termites (Isoptera) Using a Radioactive Tracer. *An Entomol Soc Am* 73(6):708-714

371 Su NY, Ye WM, Ripa R, Scheffrahn RH, Giblin-Davis RM (2006). Identification of Chilean *Reticulitermes*  
372 (*Isoptera: Rhinotermitidae*) inferred from three mitochondrial gene DNA sequences and soldier  
373 morphology. *An Entomol Soc Am* 99(2): 352-363

374 Traniello JFA, Leuthold RH (2000). Behavior and Ecology of Foraging in Termites. In: Abe Takuya, Bignell  
375 David Edward, Higashi Masahiko, editors. *Termites: Evolution, Sociality, Symbioses, Ecology*.  
376 Dordrecht: Springer Netherlands. Kluwer Academic Publishers; 2000. p. 141–68

377 Tschinkel WR (1999). Sociometry and sociogenesis of colony-level attributes of the Florida harvester ant  
378 (*Hymenoptera: Formicidae*). *An Entomol Soc Am* 92(1):80-89

379 Tschinkel WR (2010). Methods for casting subterranean ant nests. *J Insect Sc* 10(1)

380 Tschinkel WR (2011). Back to basics: sociometry and sociogenesis of ant societies (*Hymenoptera: Formicidae*).  
381 *Myrmecol News* 14:49-54

382 Turner JS (2001). On the mound of *Macrotermes michaelseni* as an organ of respiratory gas exchange. *Phys*  
383 *Biochem Zool* 74(6):798-822

384 Wenzel JW (1990). Nest design and secondary functions of social insect architecture. In G.Veeresh, B. Mallick,  
385 C. Viraktamath (Eds.), *Social insects and the environment* (pp. 657–658).  
386  
387

388 **Legends:**

389 **Figure 1.** Two-dimensional DICOM scans of *Reticulitermes* nests. Locations in which wood had been consumed  
390 (regions of interest: ROIs) were traced manually (dotted line below). Their areas were then calculated (white  
391 shapes below).

392

393 **Figure 2.** On the left is a three-dimensional image showing the exterior tunnels performed by *R. flavipes*  
394 termites. In the center and on the right are two-dimensional images of interior structures (red circle: cap; orange  
395 oval: wall; yellow circle: pillar) made with sand. Softer wood (i.e., of lower density) is black or gray, while  
396 harder wood (i.e., of higher density) is white.

397

398 **Figure 3.** Three-dimensional images of *R. grassei* nests, where sand-based and wood-based structures can be  
399 distinguished (yellow vs. brown, respectively). Here, areas in which wood is present (i.e., based on density  
400 values) have been artificially removed to allow the chambers (white) to be more easily observed.

401

402 **Figure 4.** Percentage of attacks at different locations on the wood blocks for the two termite species. The  
403 percentages were calculated using all the observations from all the colonies. The “outer sides” were the exposed  
404 faces of the block. The “underside” was the face in contact with the ground. “Interior” indicates chambers built  
405 from an earlier chamber. “After tunneling” indicates attacks initiated from within a sand tunnel that targeted the  
406 sides of the block.

407

408 **Figure 5.** Species-specific distributions of ROI volume

409

410 **Figure 6.** Boxplots of chamber number (light gray bars), volume of wood consumed (dark gray bars), and the  
411 percentage of the total wood consumed (dark line) over time in nests built by *R. grassei* (upper graph) and *R.*  
412 *flavipes* (lower graph). Within the boxes, the solid white lines are the means and the solid black lines are the  
413 medians.

414

415 **Figure 7.** Relationship between chamber number and the percentage of wood consumed for *R. grassei* (gray)  
416 and *R. flavipes* (black). The solid lines represent the best-fit regression lines, and the dotted lines indicate the  
417 95% confidence intervals.

418


419 **Figure 8.** Number of interior structures built by *R. flavipes* over time (means  $\pm$  SE). The solid black line is the  
420 best-fit regression line, and the dotted gray lines indicate the 95% confidence interval.

421

422 **Figure 9.** Relationship between chamber number and interior structure number for *R. flavipes* (means  $\pm$  SE). The  
423 solid black line is the best-fit regression line, and the dotted gray lines indicate the 95% confidence interval.

424


425 Figure 1



426

427


428 Figure 2



429


430

431 Figure 3



432

433




435

436

437


438 Figure 5



439


440

441




443

444



446

447



449

450

451






Figure 1




Figure 2.

